

How much weigh a pound in gravitational field

A.I.Nikishov *

*I.E.Tamm Department of Theoretical Physics,
P.N.Lebedev Physical Institute, Moscow, Russia*

January 28, 2014

Abstract

The force in the equation of motion of a particle should be in accordance with energy conservation in a constant gravitational field. It turns out that this is possible only if the force is given by the change of momentum per unit of coordinate (not proper) time. We discuss the consequences of this fact. In particular it turns out that the spring balance, which keeps the body at rest in (strong) gravitation field, shows the finite value even at infinitely close approach to horizons of black holes, if they exist in Nature.

There are several ways of introducing force in the equation of motion of a particle in gravitational field. So Möller elaborates three forces "to describe all aspects of nongravitational interaction", see §10.4 in [1]. Landau and Lifshitz define the gravitational force as 3-space covariant derivative of momentum over (synchronized) proper time, see §88, Problem 1 in [2]. Not all of these forces respect the energy conservation law. Those which do it are not covariant. We consider some of these approaches.

In a constant gravitational field the covariant component of momentum p_0 is conserved: (see §88 in [2])

$$p_0 = \frac{mc^2 \sqrt{g_{00}}}{\sqrt{1 - \frac{v^2}{c^2}}}, \quad v^\alpha = \frac{cdx^\alpha}{\sqrt{g_{00}}(dx^\alpha - g_{\alpha} dx^\alpha)}, \quad g_\alpha = -\frac{g_{0\alpha}}{g_{00}}, \quad v^2 = \gamma_{\alpha\beta} v^\alpha v^\beta. \quad (1)$$

For particle at rest

$$p_0 = mc^2 \sqrt{g_{00}}, \quad dp_0 = \frac{mc^2}{2\sqrt{g_{00}}} g_{00,\alpha} dx^\alpha. \quad (2)$$

Hence the coordinate nongravitational force, which adiabatically changes p_0 is

$$F_\alpha^{n.g.} = \frac{dp_0}{dx^\alpha} = \frac{mc^2}{2\sqrt{g_{00}}} g_{00,\alpha}. \quad (3)$$

When only space coordinates are transformed, F_α is 3-vector and dp_0 is an invariant and as such it can be integrated, see equation (10.116) in [1].

*E-mail: nikishov@lpi.ru

Next we assume that in a small region of space the gravitational force is directed against the axis 1. Then only dx^1 is nonzero:

$$dp_0 = \frac{mc^2}{2\sqrt{g_{00}}}g_{00,1}dx^1 = \frac{mc^2}{2}\frac{g_{00,1}}{\sqrt{|g_{00}g_{11}|}}dl^1, \quad dl^1 = \sqrt{g_{11}}dx^1. \quad (4)$$

From here it is natural to assume that the 3-invariant nongravitational force, holding particle at rest in constant gravitational field, is the force measured by a spring balance:

$$F = \frac{mc^2g_{00,1}}{2\sqrt{|g_{00}g_{11}|}}. \quad (5)$$

According to heuristic approach to gravity [3] one may expect that $|g_{00}g_{11}| = 1$. In general relativity this is true only in linear approximation in isotropic coordinates in Schwarzschild solution, which has the form

$$ds^2 = g_{00}(dx^0)^2 + g_{11}[(dx^1)^2 + (dx^2)^2 + (dx^3)^2]. \quad (6)$$

In any case $|g_{00}g_{11}|$ is always finite. Then it follows from (6) that the invariant force acting on particle at rest remains finite even when the particle is approaching the black hole horizon. This remains true also when $g_{0\alpha}$ are nonzero because the Coriolis-type force is not acting on particle at rest.

On the other hand according to eq. (2.2.6) in [4] when $v^\alpha = 0$ the gravitational force (and nongravitational one holding the particle at rest) is

$$K = m\sqrt{\gamma_{\alpha\beta}a^\alpha a^\beta}, \quad a^\alpha = \frac{d^2x^\alpha}{ds^2}, \quad \alpha = 1, 2, 3. \quad (7)$$

For the metric (6) we find

$$K = \frac{mc^2}{2}\frac{g_{00,\alpha}}{\sqrt{|g_{00}g_{11}|}}\frac{1}{\sqrt{g_{00}}} \quad (8)$$

Contrary to (5) this quantity grows unlimitedly at the approach to black hole horizon. The contradiction with (5) disappears if the derivative of momentum over the proper time is replaced by the derivative over coordinate time.

The derivative over coordinate time naturally appears in Lagrange formalism, see §10.4 in [1]. Indeed, using

$$L = -mc(g_{ik}\frac{dx^i}{dt}\frac{dx^k}{dt})^{1/2}, \quad (9)$$

we have

$$p_\mu = \frac{\partial L}{\partial \tilde{u}^\mu} = -\frac{mcg_{\mu l}\tilde{u}^l}{\sqrt{g_{ik}\tilde{u}^i\tilde{u}^k}}, \quad \tilde{u}^k = \frac{dx^k}{dt}, \quad \frac{\partial L}{\partial x^\mu} = -\frac{mcg_{lm,\mu}\tilde{u}^l\tilde{u}^m}{2\sqrt{g_{ik}\tilde{u}^i\tilde{u}^k}}. \quad (10)$$

Then the Euler- Lagrange equations are

$$\frac{dp_\mu}{dt} = \frac{\partial L}{\partial x^\mu} = -\frac{mcg_{lm,\mu}\tilde{u}^l\tilde{u}^m}{2\sqrt{g_{ik}\tilde{u}^i\tilde{u}^k}} = \overset{g.}{F}_\mu. \quad (11)$$

For particle at rest $\tilde{u}^i = c\delta_0^i$ and $\overset{g.}{F}$ agrees with (3). (for slowly moving particle $\overset{g.}{F}_\mu \approx -\overset{n.g.}{F}_\mu$). Introducing a nongravitational force we have

$$\frac{dp_\mu}{dt} = \overset{g.}{F}_\mu + \overset{n.g.}{F}_\mu. \quad (12)$$

Now we dwell briefly on Problem 1 in §88 in [2]. In the second edition of this book (Moscow, 1948) the problem is treated for small values of velocity. The force is obtained in the form

$$f_\alpha = \frac{dp_\alpha}{d\tau} = mc^2 \left[-\frac{1}{2} \frac{g_{00,\alpha}}{g_{00}} + \sqrt{g_{00}} (g_{\beta,\alpha} - g_{\alpha,\beta}) \frac{v^\beta}{c} \right], \quad g_\alpha = -\frac{g_{0\alpha}}{g_{00}}. \quad (13)$$

Replacing the differentiation over τ by differentiation over t i.e. multiplying both sides of (13) by $\sqrt{g_{00}}$ we get

$$\sqrt{g_{00}} f_\alpha = \dot{F}_\alpha = mc^2 \left[-\frac{1}{2} \frac{g_{00,\alpha}}{\sqrt{g_{00}}} + g_{00} (g_{\beta,\alpha} - g_{\alpha,\beta}) \frac{v^\beta}{c} \right]. \quad (14)$$

For particle at rest this agrees with (3). For arbitrary velocity the equation (3) in Problem 1 in §88 in [2] should agree with equations (10.75)-(10.77) in [1] up to a common factor, but this is not seen.

It is worthwhile to note that by itself the equivalence principal and the requirement of general covariance do not give the equation (3) or (5), which gives the weight of a body in constant gravitational field. So it is not a sin to try to find an algorithm for the metric, which is different from Einstein equations.

In conclusion I thank V.I.Ritus and M.I.Zelnikov for stimulating discussions.

1 References

- 1.C. Möller, *The Theory of Relativity*, Clarendon Press, Oxford (1972).
- 2.L.D.Landau and E.M.Lifshitz, *The classical theory of fields*, Moscow, (1973) (in Russian).
3. H.Dehnen, H.Hönl, and K.Westpfahl, Ann. der Phys. **6**, 7 Folge, Band 6, Heft 7-8, S.670 (1960).
- 4.I.D.Novikov, V.P.Frolov, *Physics of black holes*, Moscow (1986) (in Russian).